



**ICAR-NATIONAL RESEARCH CENTRE FOR GRAPES,
Manjri, Pune.
WEATHER DATA FOR THE PREVAILING WEEK**



Thursday (25/07/2024)– Wednesday (31/07/2024)

Location	Temperature (°C)		Possibility of Rain	Cloud Cover	Wind Speed (Km/hr) Min- Max	R H%	
	Min	Max				Min	Max
Nashik	22-23	26-29	Vani, Loni, Nashik,Dindori,Ozar, Pimpalgaon Baswant, Palkhed, Kalwan —Thu – Wed – Light to Moderate Rain .	Clear to cloudy	26-33	79-92	92-96
Pune	20-21	23-26	Pune, Phursungi, Loni Kalbhor, Uruli Kanchan, Patas, Yavat, Narayangaon,- Baramati, .Indapur –Thu – Wed –Drzzling to Light Rain .	Clear to cloudy	23-33	76-84	86-91
Solapur	21-23	28-33	Tuljapur. Latur, AUSA, Solapur .Pandharpur Vairag, Barshi, Nannaj–Thu – Wed –Drzzling to Light Rain .	Clear to cloudy	28-35	54-73	79-87
Sangli	21-22	25-28	Palsi , Khanapur Vita. Shetphal, ,Shirguppi, Walva, Palus, Miraj, Kawthe —Thu – Wed – Light to Moderate Rain .	Clear to cloudy	29-40	76-83	89-93
Vijayapura	21-22	28-31	Vijayapura, Chadchan, Tikota , Telsang– Thu – Wed –Drzzling to Light Rain .	Clear to cloudy	34-42	62-71	82-88
Hyderabad	21-23	26-30	Hyderabad, Medchal,. Zahirabad-Thu – Wed –Drzzling to Light Rain .	Clear to cloudy	26-33	56-78	80-85
Satara	20-21	24-27	Satara, Khatav, Phaltan – Thu – Wed – Drzzling to Light Rain .	Clear to cloudy	23-38	78-89	90-95
Ahmednagar	21-23	27-30	Rahata, Kopargaon, Sangamner, Akole, Shrigonda, Ahmednagar ,Karjat, Jamkhed— Thu –Wed – Drzzling to Light Rain.	Clear to cloudy	29-43	65-77	84-93
Jalna	21-23	24-31	Mantha, - Ambad, Ghansavangi, Jalna, Jafrabad– Thu – Wed – Drzzling to Light Rain .	Clear to cloudy	22-30	61-86	86-92
Buldhana	22-23	25-30	Chikhli Sindkhedraja, D.raja , Buldana – Thu – Wed –Drzzling to Light Rain .	Clear to cloudy	21-26	74-86	88-92

Kolhapur	22-23	27-28	Kagal, Karveer, Gagan-bavada – Thu -Wed – – Light to Moderate Rain.	Clear to cloudy	14-25	88-92	93-96
Bengaluru Rural	20-21	27-28	Anekal, Doddaballapur, Bengaluru-east, Bengaluru-north, Bengaluru- - Thu - Wed - Drzzling to Light Rain .	Clear to cloudy	25-32	55-61	77-84
Belagavi	21-22	25-26	Belagavi, Gokak , Chikodi, Athni - Thu - Wed Drzzling to Light Rain .	Clear to cloudy	20-30	87-91	91-95
Bidar	21-22	26-30	Basavakalyan, Humanabad, Bidar— Thu – Wed – Drzzling to Light Rain .	Clear to cloudy	22-34	64-83	86-91
Bagalkot	21-22	27-30	Hungund , Bagalkot , Jamkhandi, Mudhol –Thu – Wed – Drzzling to Light Rain .	Clear to cloudy	29-39	61-74	80-87

Note: Above weather information is summary of weather forecasting given in following websites

https://www.wunderground.com/?cm_ven=cgi

<https://imdagrmet.gov.in/weatherdata/BlockWindow.php>

<https://www.timeanddate.com/weather/india>

ICAR-National Research Centre for Grapes does not claim accuracy of it.

II. Water management

a. Number of days after foundation pruning: 93

b. Expected Pan evaporation: 0 to 2.0 mm

Amount of irrigation advised :

1. All the grape growing regions are forecasted to receive from drizzling to moderate rains. In case the soil is under wapsa (field capacity) condition, do not irrigate the vineyard.
2. If the soils especially medium and heavy, are saturated with water, then, do not irrigate for atleast 5-7 days till the soil comes to wapsa condition.
3. During fruit bud differentiation stage, shoot vigour to be controlled and hence, the irrigation water applied should be around 1000 L/ acre/ day.
4. **Cane maturity stage:** Apply irrigation through surface drip @ 1000 L/acre per day.

5. In case of monsoon rains, remove mulch cover on the bund and allow the rain water to seep into the soil. This will leach the accumulated salts in the rootzone. The mulch so removed can be mixed with the soil to improve the soil porosity.

Soil and Nutrient management :

1. In many of the grape growing areas, continuous spells of rains have been received and further also possibility of rains are there. The soils are already saturated. This has affected the rooting activity. Due to prolonged saturation, the roots may have started decaying. **Donot disturb the soil in the root zone. Wait for the soil to come to the wapsa condition before any soil related intervention has to be done.** Growth will be slow and cane maturity will be affected but donot worry. Only after wapsa, fertilizer application should be done.
2. Due to continuous sprays the leaf will not look healthy, need based sprays should be followed as the leaf health is bound to affect the photosynthate formation. This will impact cane maturity.
3. After current rains, give foliar spray of SOP @ 3-5 g/L depending upon canopy size.
4. In case of calcareous soils where acute iron deficiency is observed, repeatedly spray 2-3g/L Ferrous sulphate two to three times at 3 days interval followed by 15-20 kg/ acre Ferrous sulphate application through drip. The fertigation dose should be split into atleast 3 doses of 5kg each. Apply 5kg/ acre soluble sulphur through drip every week. Also spray magnesium sulphate and potassium sulphate @ 3 gm each/ L once only.
5. Possibility of leaf curling, check the leaf margins, if slight to more yellow, possibility of potassium deficiency. Foliar spray of SOP @ 3-4g/L followed by fertigation of 20-25 kg SOP/acre in 2 to 3 splits.
6. In coloured varieties like Jumbo, Nanasaheb Purple etc., leaf curling along with reddening/ bronzing of the leaf margin can be observed if potassium deficiency is there. Foliar spray of SOP @ 3g/L followed by fertigation of 20-25 kg SOP/acre in 2 to 3 splits.

Cane maturity stage

1. After current rains, give foliar spray of SOP @ 4-5 g/L depending upon canopy.
2. Potassium application is required from Cane maturity stage onwards. Approx. 64 kg of sulphate of potash (soluble grade) should be applied in this stage. Split the application into atleast five doses to reduce the leaching losses of the potassium. Apply 15 kg SOP in two – three splits during this week. In calcareous soils, provide foliar application of Sulphate of Potash (@ 4g/L) once in this growth stage.
3. Apply magnesium sulphate @ 15 kg/acre in two splits. The application should be done during 60-75 days after pruning. In calcareous soils, provide foliar application of Magnesium sulphate (@3g/L) in this growth stage.
4. In case of calcareous soils where acute iron deficiency is observed, repeatedly spray 2-3g/L Ferrous sulphate two to three times at 3 days interval followed by 15-20 kg/acre Ferrous sulphate application through drip. The fertigation dose should be split into atleast 3 doses of 5kg each.
5. To effectively manage calcareous soil, apply 5kg/ acre soluble sulphur through drip every week. Also spray magnesium sulphate and potassium sulphate @ 3 gm each/ L once only.
6. In case due to rains and for preventive control, if bordeaux or copper sprays are given, then there is possibility of leaf reddening in coloured varieties like Krishna Seedless etc. No specific pattern will be there. This may be due to copper toxicity. Regulate copper sprays.
7. After cane maturity, raise Sunnhemp or Dhaincha for green manuring purpose.

Pre-pruning operations – Fruit pruning season

1. In case pruning is planned during August - September, raise Sunnhemp or Dhaincha for green manuring purpose.
2. The vineyards where sodicity problems are there, apply gypsum to the soil for removal of sodium from the soil exchange complex. In case of calcareous soils, use sulphur for similar purpose. The application should be alongwith FYM/compost etc. They should be mixed in the soil and not left on the top.

3. In case of calcareous soils, if SSP is applied as basal dose, mix with FYM/compost etc. to avoid phosphorus fixation.
4. Test the soil and irrigation water, to plan for nutrient and water management during fruit pruning season.
5. In areas where rains have not been received and the irrigation water availability is less, it is suggested to flood the rootzone(only) with water to leach out the salts and wet the entire soil depth before pruning and then cover with mulch. Thereafter irrigate as per availability of water.

III. Canopy Management

Based on the weather data and growth stages, following suggestions are offered for vineyard management.

A) Old vineyard:

1. The recent rainfall has increased humidity in the atmosphere. This will help to increase the shoot vigour.
2. The new growth coinciding with continuous rains will make the vine more susceptible for fungal diseases like downy mildew and anthracnose.
3. The vineyard after 90 days might be in the stage of cane maturity. If the new growth continues, the cane maturity will be delayed. Shoot pinching and removal of side shoot will help to create open canopy thereby reducing the chances of humidity build-up. The disease load will be reduced with effective coverage.
4. Training of shoots on foliage wire will help for aeration in the canopy thereby reducing the chances of diseases. This will also help for uniform spray coverage of insecticides/ fungicides used for the control of pest and diseases.
5. Application of potash through drip @ 1.0 to 1.25 kg per acre and spray @ 3.5 to 4.0 g/L water will help to advance cane maturity.
6. In many vineyards, irregular cane maturity is experienced by the grape growers. Bourdeaux spray @ 0.75 to 1.0% at 10 days interval will help to control the problem.

7. In case of leaf yellowing which is basically due to the deficiency of ferrous, magnesium and potash created by availability of calcium carbonate in excess amount in the soil. Hence, application of sulphur mixed with FYM will help to overcome the problem. At this stage, FYM application may not be possible. Under such condition, soluble grade ferrous and magnesium can be applied through drips.
8. To achieve regular cane maturity, application of phosphorous and potash grade fertilizer can be applied in the soil. Avoiding the stress to the vine will control this problem to certain extent. Spraying of Boudreaux mixture @ 0.75 to 1.0% and also drenching through soil will help to control.

B) New vineyard:

- 1) Due to recent rains in vineyard, there will be vigorous growth of new shoots. Under the situation of delayed cordon development, spraying of cytokinin based PGR (6BA @ 10 ppm) will help to increase cytokinin and reduce gibberellin level in the vine.
- 2) Control of shoot vigour is most important to achieve fruit bud differentiation. Hence, potash to be sprayed at minimum concentration of 2.0 to 2.5 g/L water.
- 3) Considering the bud differentiation, 2 to 3 sprays of 0.52.34 @ 2.0 to 2.5 g/L water can be given.
- 4) Extension of cordon will help to develop fruitful canes. At this stage, application of soluble fertilizer like 12:61:0 @ 1.25 to 1.5 kg/acre should be applied. In addition, DAP @ 25kg/acre as a basal dose should also be applied.
- 5) Since the shoot growth is coinciding with rainfall and high humidity, application of cytokinin based PGR will help to accelerate fruit bud differentiation. Once the new shoots are of 5-6 leaf stage, pinching should be done at 4-5 leaf. This should be followed by spraying of 0:52:34 @ 2.0 to 2.5 g/L water.

C) Rootstock plots:

- 1) The period of grafting new varieties on rootstock is approaching. Selection of specific scion is more important considering the life span of any variety.
- 2) Preparation of rootstock 10 days before grafting is required. Retention of 3-4 straight growing, vigorous and healthy rootstock shoots should be done before the grafting. In case of excess shoots available, shoot thinning to be done. In addition, the removal of side shoots

in at least two instalments can be done to achieve straight and thick shoot (approx. 8.0 mm) at 1.0 feet height above the ground.

IV. Disease management

Days after foundation pruning	Risk of diseases			
	Downy mildew	Powdery mildew	Anthracnose	Others (specify)
93	Low	Low	Moderate	Bacterial spot- High Rust-Nil

Two sprays of Kasugamycin 5% +Copper Oxychloride 45% WP @750g/ha, may be given in all grape growing areas to manage bacterial spot and anthracnose due to the continuous rains presently in most of the areas. Application of Thiophenate methyl/carbendazim @1g/L will provide a good control against anthracnose. Downy mildew can be prevented by application of mancozeb which can also control bacterial spot. In Sangli area where powdery mildew attacks are reported, an application of triazoles like hexaconazole @1ml/L might help. A foliar application of Trichoderma@ 4-5ml/L may be given as the moisture conditions will be suitable for multiplication of the biocontrol agents. Drip application of Trichoderma should continue at fortnightly intervals but it can be put in hold till the rain stops in Sangli and adjoining regions.



Bacterial spot



Anthracnose

V. Insect and Mite management

1. Due to optimum weather conditions, mealybug infestation may be noticed. Use of broad-spectrum insecticides should be avoided for mealybug control. Preventive plant wash, on stem and cordons, of biocontrol agents such as *Verticillium*, *Metarhizium*, *Beauveria* may be given. In case of shoot malformation, remove excess shoot growth and give foliar spray of imidacloprid 17.8 SL @ 0.4 ml per litre water.

2. In case of thrips or caterpillar infestation, remove excess canopy. Application of fipronil 80 WG @ 0.0625 g per litre or emamectin benzoate 5 SG @ 0.22 g per litre water is effective. Light traps may be installed outside the vineyards to manage moths for reducing caterpillar infestation.
3. Mite infestation may start appearing, therefore, monitor the vineyards carefully. If mite infestation is observed, sulphur 80 WDG @ 1.5-2.0 gram per litre or abamectin 1.9 EC @ 0.75 ml/l water is effective.
4. Red colour stem borer (*Dervishiya cadambae*) has started egg laying and infestation under bark in grape areas. Install light traps near the vineyards to manage moths of this stem borer. Remove loose bark from stem and cordons and give preventive wash on stem and cordons with biocontrol agent *Metarhizium* @ 3-5 ml per litre water minimum once in the month during July to September months. If infestation is observed, remove the loose bark and give spot stem and cordon wash with *Metarhizium* @ 3-5 ml per litre water and 1.5-2 litres water per plant on the infested plants only.
5. In new vineyards, flea beetle infestation may be observed. In case of heavy infestation, give soil drenching with imidacloprid 17.8 SL @ 1.5 ml per plant and foliar application with spinetoram 11.7 SC @ 0.3 ml per litre or fipronil 80 WG @ 0.0625 g per litre water.

CONFIDENTIAL